

ADA08 - 9am Thu 29 Sep 2022

Parameter Uncertainties
Confidence Intervals and Regions

Fitting a Linear Trend
Orthogonal vs Correlated Parameters

Summary of the ADA Roadmap:

Algebra of Random Variables


Minimising BoF = χ^2

Alternative (Robust) BoFs

Maximum Likelihood (ML) BoF=- $2\ln L$

Bayesian Inference (MAP) BoF=- $2\ln(L P)$

Max Likelihood and Bayesian Inference




Monte-Carlo Error Propagation

1. Create mock datasets.

1a. “Jiggle” the data points (using Gaussian random numbers).

* Requires good error bars.



1b. (and/or) “Bootstrap” samples:

Pick N data points at random, with replacement (some points omitted, some repeated).


* Requires more data than parameters ($N > M$).

* Works with no error bars available.

2. Fit the model to each mock dataset.

$$\langle X_i \rangle = a t_i + b$$

3. Observe how the best-fit parameter values “dance”.



4. Accumulate histograms approximating the parameter probability distributions.

5. Compute mean, median, variance, MAD, etc. of the parameters, or **any function of the parameters**.

Confidence interval on a single parameter

(1-parameter, k-sigma confidence interval)

The **1- σ confidence interval** on α includes 68% of the area under the likelihood function:

$$L(\alpha) \equiv P(X | \alpha) \propto \frac{e^{-\chi^2/2}}{\prod_i \sigma_i}$$

or posterior probability distribution, for non-uniform prior $P(\alpha)$:

$$P(\alpha | X) \propto L(\alpha) P(\alpha)$$


For a **k - σ (1-parameter) confidence interval**, use $\Delta\chi^2 = k^2$,

$\Delta\chi^2 = 1$ for 1- σ , 68% probability

$\Delta\chi^2 = 4$ for 2- σ , 95.4% probability

$\Delta\chi^2 = 9$ for 3- σ , 99.73% probability


...



Generalise:

$$\chi^2 \Rightarrow -2 \ln(L(\alpha) P(\alpha))$$

2-parameter 1-sigma Confidence Region



If Y is a “nuisance parameter”, use the **1-parameter 1-sigma confidence interval** in X , *tangent to* the $\Delta\chi^2 = 1$ contour in (X, Y) .
This interval encloses 68% probability.

If both X and Y are of interest, use the **2-parameter 1-sigma confidence region**, the $\Delta\chi^2 = 2.30$ contour in (X, Y) .
This contour encloses 68% probability.

Use $-2 \ln(L(\alpha) P(\alpha))$ instead of χ^2 , if needed.

Note: Contour enclosing 68% probability must be **wider** than the 1-parameter confidence interval.


Why?

$$L(\alpha) \equiv P(X | \alpha) \propto \frac{e^{-\chi^2/2}}{\prod_i \sigma_i}$$

M -parameter $k\sigma$ Confidence Regions

$\Delta\chi^2$ thresholds for M -parameter $k\sigma$ Confidence Regions

	Prob	$M = 1$	2	3	4
$1 - \sigma$	68%	1	2.30	3.53	4.72
$2 - \sigma$	95.4%	4	6.17	8.02	9.70
$3 - \sigma$	99.73%	9	11.8	14.2	16.3



The **M -parameter confidence region** is enclosed by the $\Delta\chi^2$ surface including the desired probability.

All **nuisance parameters must be re-fitted** (or integrated over) for each set of fixed values for the M parameters in the sub-space of interest.

(a.k.a. “marginalise over the nuisance parameters”.)

The $\Delta\chi^2$ in the M -parameter sub-space has a χ^2_M distribution, with M degrees of freedom.

Example: Estimate both μ and σ

$$L(\mu, \sigma) \equiv P(X | \mu, \sigma) = \frac{e^{-\chi^2/2}}{(2\pi)^{N/2} \sigma^N}$$

$$-2 \ln L = \sum_{i=1}^N \left(\frac{X_i - \mu}{\sigma} \right)^2 + 2N \ln \sigma + \text{const}$$


$$0 = \frac{\partial}{\partial \mu} [-2 \ln L] = -2 \sum_{i=1}^N \frac{X_i - \mu}{\sigma^2}$$

$$0 = \frac{\partial}{\partial \sigma} [-2 \ln L] = -2 \sum_{i=1}^N \frac{(X_i - \mu)^2}{\sigma^3} + \frac{2N}{\sigma}$$


$$\mu_{\text{ML}} = \frac{1}{N} \sum_i X_i \quad \sigma_{\text{ML}}^2 = \frac{1}{N} \sum_i (X_i - \mu_{\text{ML}})^2$$

Posterior \propto Likelihood \times Prior

$$P(\mu, \sigma | X) \propto L(\mu, \sigma) P(\mu, \sigma)$$




2-parameter 1,2,3- σ confidence regions:
 $L(\mu, \sigma) = P(X|\mu, \sigma)$




Note: ML gives biased estimate for σ .


Example: Estimate both μ and σ




$$L(\mu, \sigma) = P(X|\mu, \sigma)$$



$$L(\mu, \sigma) = P(X|\mu, \sigma)$$





$$L(\mu, \sigma) = P(X|\mu, \sigma)$$



Contours: 1,2,3-sigma 2-parameter confidence regions for μ and σ .

Dashed curves: maximum-likelihood estimates for μ_{ML} and σ_{ML} .

True values: $\mu=0$ and $\sigma=1$.



Fit a line to $N=1$ data point

Fit $y = a x + b$ to $N=1$ data point:


Blue lines : $\chi^2 = 0$

Red lines : $\chi^2 = 1$

χ^2 contours in the (a,b) plane:



Solution is **degenerate**, since $M=2$ parameters are constrained by only $N=1$ data point.

Bayes: prior $P(a,b)$ needed to determine a unique solution.




Fit a line to $N = 2$ data points

- Fit $y = a x + b$ to $N = 2$ data points:
 - red lines give $\chi^2 = 2$
 - blue line gives $\chi^2 = 0$
- Note that a, b are not independent.





Correlated Parameters 😞

- Parameters a and b are correlated : (
- To find the optimal (a, b) we must:
 - minimize χ^2 with respect to a at a sequence of fixed b values
 - then minimise the resulting χ^2 values with respect to b .
- If a and b were independent, then all slices through the χ^2 surface at each fixed b would have same shape and minimum.
- Similarly for a .
- We could then optimize a and b independently, saving a lot of calculation
- **How to make a and b independent of each other?**



Orthogonal Parameters for fitting a line to $N = 2$ data points

- Fit $y = a(x - \hat{x}) + b$
- Different parameters for same model.
- Note: a, b are now independent !






Orthogonal slope and intercept


Analysis using the algebra of random variables:

$$y = a(x - \hat{x}) + b$$

$$\hat{x} = \frac{x_1 + x_2}{2}$$




Corresponds to $\Delta\chi^2 = 1$.




Orthogonal vs Correlated Parameters

Correlated Parameters ☹



Orthogonal Parameters 😊



For each a , a different b minimises χ^2 .
For each b , a different a minimises χ^2 .

For any a , the same b minimises χ^2 .
For any b , the same a minimises χ^2 .

Fit a line to N data points

- If we use $y = a x + b$
then a, b are correlated.
- Make a, b orthogonal:

$$y = a(x - \hat{x}) + b$$

$$\hat{x} = \frac{\sum x_i / \sigma_i^2}{\sum 1 / \sigma_i^2} \quad (\text{to be proven later})$$

- Intercept: Set $a = 0$ and optimise b :


optimal average:

$$\hat{b} = \hat{y} = \frac{\sum y_i / \sigma_i^2}{\sum 1 / \sigma_i^2}, \quad \text{Var}[\hat{b}] = \frac{1}{\sum 1 / \sigma_i^2}$$

- Slope: Set $b = 0$ and optimise a :


optimal scaling of pattern: $P_i = x_i - \hat{x}$

$$\hat{a} = \frac{\sum y_i (x_i - \hat{x}) / \sigma_i^2}{\sum (x_i - \hat{x})^2 / \sigma_i^2}, \quad \text{Var}[\hat{a}] = \frac{1}{\sum (x_i - \hat{x})^2 / \sigma_i^2}$$



Choose Orthogonal Parameters

- Good practice (when possible).
- Results for any one parameter don't depend on values of other parameters.
- Example: fit a gaussian profile.
2 fit parameters:
 - Width, w
 - Area or peak value. Which is best?



Peak value depends
on width – bad

$$f(x) = Pe^{-\frac{1}{2}\left(\frac{x-x_0}{w}\right)^2}$$

$$g(x) = \frac{A}{w\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-x_0}{w}\right)^2}$$

Area is (more nearly)
independent
of width – good

Fini -- ADA 08