

Lecture 14


Early Inflation

A possible solution for:


Why a hot Big Bang?
 Horizon problem
 Flatness problem
 Monopole problem
 Origin of primordial structure

AS 4022 Cosmology

Early inflation, Hot Big Bang, Decelerating Expansion, Late inflation by "Dark Energy"



Early Inflation: the first 10^{-33} seconds



1980: Inflation (Alan Guth)

- Universe born from "nothing" ?
- A **quantum fluctuation** produces a tiny bubble of "False Vacuum".
- High vacuum energy drives **exponential expansion**, also known as "inflation."
- Universe expands by huge factor in tiny fraction of second, as false vacuum returns to true vacuum.
- Expansion so fast that **virtual particle-antiparticle pairs** get separated to become **real particles and anti-particles**.
- Stretches out all structures, giving a **flat geometry** and uniform T and ρ , with **tiny ripples**.
- Inflation launches the **Hot Big Bang!**

AS 4022 Cosmology

Planck Units

Planck Length: de Broglie wavelength \sim Schwarzschild radius

$$E = Mc^2 = \frac{hc}{\lambda} \Rightarrow \lambda = \frac{h}{Mc} \quad R_s = \frac{2GM}{c^2}$$

$$(\lambda R_s)^{1/2} \sim L_p = \left(\frac{\hbar G}{c^3} \right)^{1/2} \sim 10^{-35} \text{ m}$$

Planck Time $t_p = \frac{L_p}{c} = \left(\frac{\hbar G}{c^5} \right)^{1/2} \sim 10^{-43} \text{ s}$

Planck Mass $M_p = \frac{L_p c^2}{G} = \frac{\hbar}{L_p c} = \left(\frac{\hbar c}{G} \right)^{1/2} \sim 10^{25} m_p \sim 10^{19} \text{ GeV}/c^2$

Planck Energy $E_p = M_p c^2 = \left(\frac{\hbar c^5}{G} \right)^{1/2} \sim 10^{19} \text{ GeV}$

Limits of Quantum Mechanics and General Relativity. Need Quantum Gravity theory (as yet unknown) to describe physics at these scales.

AS 4022 Cosmology

Quantum Vacuum


Heisenberg Uncertainty principle:

$$\Delta E \Delta t \sim \hbar$$

Can violate energy conservation:

"Borrow" energy from the vacuum, but only for a short time.

To create a particle-antiparticle pair, need $\Delta E > 2 m c^2$. These "virtual" pairs live only briefly, $\Delta t \sim \hbar / \Delta E$.



Vacuum is not empty. Filled with all types of virtual pairs.

AS 4022 Cosmology

Quantum Vacuum

Like waves on the sea.

Quantum fields oscillate in many possible wave modes

$$\phi(\mathbf{x}, t) \quad \psi_i(\mathbf{x}, t) \quad \mathbf{A}(\mathbf{x}, t), \dots$$

Each wave mode is a Harmonic Oscillator,

Potential energy:

$$V(\phi, \psi_i, \mathbf{A}, \dots) = m_\phi |\phi|^2 + m_\psi |\psi|^2 + \dots$$

Ladder of discrete Energy States:


$$E(n) = \left(n + \frac{1}{2}\right) \hbar \omega$$

Zero-point energy:

$$E(0) = \frac{1}{2} \hbar \omega = \frac{\hbar c}{2\lambda}$$


Quantum vacuum = all wave modes in ground state.

AS 4022 Cosmology



Vacuum Energy Density

Waves in a Box:



Density of states in 6-D phase space:

$$L = n\lambda \quad k = \frac{2\pi}{\lambda} = \frac{2\pi n}{L} \quad \Delta k = \frac{2\pi}{L}$$

$$\frac{dn}{d^3\mathbf{x} d^3\mathbf{k}} = \frac{g}{L_x L_y L_z \Delta k_x \Delta k_y \Delta k_z} = \frac{g}{(2\pi)^3}$$

$$\epsilon_{\text{vac}} = \iiint \langle E(\mathbf{k}) \rangle \frac{dn(\mathbf{k})}{d^3\mathbf{x} d^3\mathbf{k}} d^3\mathbf{k}$$

$$= \int_0^\infty \left(\frac{\hbar c k}{2}\right) \left(\frac{g}{(2\pi)^3}\right) 4\pi k^2 dk$$

$$= \frac{g \hbar c}{4\pi^2} \int k^3 dk = \frac{g \hbar c}{16\pi^2} (k_{\text{max}}^4 - k_{\text{min}}^4)$$

Short waves dominate.

Photons have $g = 2$ polarisations.
Electrons have $g = 2$ spin states.

Each mode is a simple harmonic oscillator.

Zero-point energy per mode:

$$\langle E(\mathbf{k}) \rangle = \frac{1}{2} \left(\frac{\hbar c}{\lambda}\right) = \frac{\hbar c k}{2}$$

AS 4022 Cosmology

Cosmological Constant Problem

Range of wavelengths in the Cosmological box:

$$\lambda > L_p \sim \left(\frac{\hbar G}{c^3}\right)^{1/2} \sim 10^{-35} \text{ m} \quad k_{\text{max}} \sim \frac{2\pi}{L_p}$$

$$\lambda < \frac{c}{H_0} \sim 4300 \text{ Mpc} \sim 10^{26} \text{ m} \quad k_{\text{min}} \sim \frac{2\pi}{c/H_0}$$

$$\epsilon_{\text{vac}} = \frac{g \hbar c}{2\pi^2} (k_{\text{max}}^4 - k_{\text{min}}^4) \approx \frac{g \hbar c}{16\pi^2} \left(\frac{2\pi}{L_p}\right)^4 = \frac{\pi^2 g \hbar c}{L_p^4} \sim \frac{E_p}{L_p^3}$$

$$\Omega_{\text{vac}} = \frac{\rho_{\text{vac}}}{\rho_{\text{crit}}} \sim \left(\frac{\pi^2 g \hbar}{c L_p^4}\right) \left(\frac{8\pi G}{3H_0^2}\right) \sim \frac{8\pi^3 g}{3} \left(\frac{c/H_0}{L_p}\right)^2 \sim 10^{120}$$

Cosmological Constant problem:

Observe $\Omega_\Lambda \sim 0.7$ Predict $\Omega_\Lambda \sim 10^{120}$

Why is Ω_Λ so small, yet not exactly zero?

Why does quantum vacuum energy not gravitate?

AS 4022 Cosmology

Flatness Problem

Why was the initial geometry flat?

If exactly flat, then always so.

To be **roughly** flat today, $\Omega_0 = 1 + \epsilon$

must initially be **incredibly** close to flat:

$$\Omega(x) = \frac{8\pi G\rho}{3H^2} = \frac{\Omega_R x^4 + \Omega_M x^3 + \Omega_\Lambda}{\Omega_R x^4 + \Omega_M x^3 + \Omega_\Lambda + (1 - \Omega_0) x^2}$$

$$\Omega(x \rightarrow \infty) - 1 \Rightarrow \frac{\epsilon}{\Omega_R x^2} \sim 10^{-119} \left(\frac{\epsilon}{0.01}\right) \left(\frac{10^{-5}}{\Omega_R}\right) \left(\frac{10^{61}}{x}\right)^2$$

$$\Omega_R \sim 10^{-5} \quad x \sim \frac{c/H_0}{L_p} \sim 10^{61}$$

Flatness problem: Why was Ω initially so close to 1?
How did the Universe know precisely how fast to expand?

AS 4022 Cosmology

Horizon Problem

Why is the universe almost perfectly isotropic?

Distant regions were never in causal contact.

Yet we see:

Same CMB temperature to 1 part in 10^5 .

Similar galaxy distributions.


AS 4022 Cosmology

The Hubble Deep Fields


Thousands of galaxies,
just a few stars.

Similar galaxy distributions,
supporting the Cosmological Principle.

HDF North




HDF South



Hubble Deep Field HST · WFPC2
NASA · ST ScI OPO · January 15, 1995 · R. Williams (ST ScI, NASA)

Hubble Deep Field South Hubble Space Telescope · WFPC2
NASA · ST ScI OPO · January 15, 1995 · R. Williams (ST ScI, NASA)



The Horizon: How far can we see?

suppose $R(t) = R_0 (t/t_0)^\alpha$

$$\chi_H = \eta_0 = \int_{t_i}^{t_0} \frac{c dt}{R(t)} = \frac{c t_0^\alpha}{R_0} \int_{t_i}^{t_0} \frac{dt}{t^\alpha} = \left(\frac{c t_0}{R_0} \right) \times \begin{cases} 1 - (t_i/t_0)^{1-\alpha} & \text{if } \alpha < 1 \\ \ln(t_0/t_i) & \text{if } \alpha = 1 \\ \frac{(t_0/t_i)^{\alpha-1} - 1}{\alpha-1} & \text{if } \alpha > 1 \end{cases}$$

As $t_i \Rightarrow 0$,
Finite for $\alpha < 1$,
Infinite for $\alpha = 1, > 1$.

matter - dominated :
 $\alpha = 2/3$
 radiation - dominated :
 $\alpha = 1/2$

AS 4022 Cosmology

Particle Horizon at $z = 1100$

$$L_H(t_R) = \int_0^{t_R} \frac{R(t_R)c dt}{R(t)} = \frac{c}{(1+z)} \int_{1+z}^{\infty} \frac{dx}{H(x)} \approx \frac{c}{(1+z)H_0} \int_{1+z}^{\infty} \frac{dx}{\sqrt{x^4 \Omega_R + x^3 \Omega_M}}$$

$$= \frac{2c}{(1+z)H_0 \sqrt{\Omega_M x_0}} \left(\sqrt{1 + \frac{x_0}{1+z}} - 1 \right) \quad x_0 = \frac{\Omega_M}{\Omega_R} \approx 3500 \left(\frac{\Omega_M}{0.3} \right)$$

$$= \frac{c}{H_0} \frac{2(\sqrt{4.6} - 1)}{1100 \cdot \sqrt{0.3} \cdot 3500} = 3.4 \times 10^{-5} \frac{c}{H_0} \approx 190 \left(\frac{0.7}{h} \right) \left(\frac{0.3}{\Omega_M} \right)^{1/2} \text{ kpc}$$

Expands by factor $1+z=1100$ to $\sim 200 \text{ Mpc}$ today.

How did these 20,000 causally disconnected regions know what temperature to be, to 1 part in 10^5 ?

AS 4022 Cosmology

Magnetic Monopole Problem

The GUT (Grand Unified Theory) phase transition should produce "topological defects" that look like magnetic monopoles.

We don't see any.

Why is the universe devoid of magnetic monopoles ?

Phase Transitions among Forces

Today: $kT \sim 10^{-36} \text{ eV}$

AS 4022 Cosmology

Inflation launches the Big Bang

expand by factor $f = e^{Ht}$ in time $t = \frac{\ln f}{H} \quad H \sim \left(\frac{8\pi G \rho_{\text{vac}}}{3} \right)^{1/2}$

What came before $t = 0$?
 Early inflation replaces the Big Bang singularity.
 Launches the Hot Big Bang.

Horizon problem: Expands a causally-connected region.
 Flatness problem: Flattens the geometry.
 Monopole problem: Moves primordial monopoles beyond horizon.
 Seeding structures: Stretches out small quantum fluctuations.

AS 4022 Cosmology

During inflation:

Particle horizon expands

$$R \propto e^{Ht} \quad L_H(t_2) = \int_{t_1}^{t_2} \frac{R(t_2)c dt}{R(t)} = \frac{c}{H} e^{H(t_2-t_1)}$$

Dramatic cooling

$$x \equiv 1+z \propto \frac{1}{R} \propto e^{-Ht} \quad T = T_0(1+z) \propto \frac{1}{R} \propto e^{-Ht}$$

Geometry flattens

$$\Omega(x) = \frac{8\pi G \rho}{3H^2} = \frac{\Omega_R x^4 + \Omega_M x^3 + \Omega_\Lambda}{\Omega_R x^4 + \Omega_M x^3 + \Omega_\Lambda + (1-\Omega_0) x^2}$$

$$\Rightarrow \frac{\Omega_\Lambda}{\Omega_\Lambda + (1-\Omega_0) x^2} \approx 1 - \frac{(1-\Omega_0)}{\Omega_\Lambda} x^2$$

AS 4022 Cosmology

How much inflation needed?

Present horizon: $R_H \sim c / H_0 \sim 4300 \text{ Mpc} \sim 10^{26} \text{ m}$

Planck time: $t_p \sim 10^{-43} \text{ s}$ horizon $\sim c t_p \sim L_p \sim 10^{-35} \text{ m}$

post-inflation redshift: $z_p \sim \frac{E_p}{k T_{\text{CMB}}} \sim \frac{10^{19} \text{ GeV}}{10^{-3.6} \text{ eV}} \sim 10^{32}$

$L_p \Rightarrow L_p f z_p \sim f \times 10^{-3} \text{ m}$ **Need inflation factor $f \sim 10^{29} \sim e^{67}$**

GUT phase transition: $E_{\text{GUT}} \sim 10^{15} \text{ GeV}$ $E \sim kT \sim R^{-1} \sim t^{-1/2}$

$t_{\text{GUT}} \sim \left(\frac{E_p}{E_{\text{GUT}}}\right)^2 t_p \sim 10^8 t_p \sim 10^{-35} \text{ s}$ horizon $\sim c t_{\text{GUT}} \sim 10^8 L_p \sim 10^{-27} \text{ m}$

redshift $\sim z_{\text{GUT}} \sim \frac{E_{\text{GUT}}}{k T_{\text{CMB}}} \sim \frac{10^{15} \text{ GeV}}{10^{-3.6} \text{ eV}} \sim 10^{38}$

$L_{\text{GUT}} \Rightarrow L_{\text{GUT}} f z_{\text{GUT}} \sim f \times 10^9 \text{ m}$

Inflation factor $f \sim 10^{17} \sim e^{40}$

Time $t \sim \ln(f) / H$
 $\sim 40 t_{\text{GUT}} \sim 10^{-33} \text{ s}$

AS 4022 Cosmology

Requirements for Inflation:

Friedmann momentum equation:

$$\frac{\ddot{R}}{R} = -\frac{4\pi G}{3c^2}(\epsilon + 3p)$$

$$= -H_0^2 \sum_w \Omega_w x^{-3(1+w)} \frac{1+3w}{2}$$

Acceleration requires a sufficiently negative pressure:

$$p < -\epsilon / 3$$

$$w < -1 / 3$$

Inflation requires the dominant component to have $w < -1 / 3$.

"Dark Energy" is driving inflation now:

$$w_\Lambda = -1 \quad p_\Lambda = -\epsilon_\Lambda$$

$$\frac{\ddot{R}}{R} \Rightarrow +\frac{8\pi G}{3c^2} \epsilon_\Lambda = H_0^2 \Omega_\Lambda$$


But today's Dark Energy is negligible at early times.

$$\Rightarrow R \propto \exp(H_\Lambda t) \quad H_\Lambda = H_0 \sqrt{\Omega_\Lambda} = \frac{\Lambda}{3}$$

AS 4022 Cosmology

What causes Early Inflation? An evolving Scalar Field?

Harmonic potential




Klein-Gordon equation
= wave equation for a massive scalar field.

$$E^2 = p^2 c^2 + m^2 c^4$$

$$E \rightarrow i \hbar \frac{\partial}{\partial t} \quad p \rightarrow -i \hbar \nabla$$

$$\ddot{\phi} - c^2 \nabla^2 \phi = \frac{m^2 c^4}{\hbar^2} \phi = \frac{m^2 \phi}{(M_p t_p)^2}$$

Higgs potential



For a spatially-uniform field:

$$\nabla \phi = 0 \quad \ddot{\phi} = -\frac{dV}{d\phi}$$


$$\text{massive field: } V = \frac{m^2 \phi^2}{2(M_p t_p)^2}$$

$$\text{Higgs field: } V \propto -a\phi^2 + b\phi^4$$

Vacuum energy starts large, declines to 0 at late times.

AS 4022 Cosmology

Scalar Field Dynamics




Kinetic energy of the oscillations is damped. Re-heats the Universe, creating all types of particle-antiparticle pairs, launching the Hot Big Bang.

AS 4022 Cosmology


Spontaneous Symmetry Breaking

Why start in the false vacuum, not near $V=0$?

Before GUT phase transition:



After GUT phase transition:



Tunnel to true vacuum? No.

Latent heat released, converts to particle-antiparticle pairs, re-heating the universe.

Small excess (10^{-9}) of particles.

AS 4022 Cosmology

Scalar Field Equation of State

Equation of State: $w = p / \epsilon$

Uniform field: $c^2 (\nabla \phi)^2 \ll \dot{\phi}^2$

$$p_\phi = \frac{1}{2} \frac{1}{\hbar c^3} (\dot{\phi}^2 + c^2 \nabla \phi^2) - V(\phi)$$


(Inflation makes spatial gradients small.)

$$\epsilon_\phi = \frac{1}{2} \frac{1}{\hbar c^3} (\dot{\phi}^2 + c^2 \nabla \phi^2) + V(\phi)$$

Required for inflation:

$$w = \frac{p}{\epsilon} = \frac{\dot{\phi}^2 - 2\hbar c^3 V}{\dot{\phi}^2 + 2\hbar c^3 V} < -\frac{1}{3}$$

$$\epsilon + 3p = 2 \left(\frac{\dot{\phi}^2}{\hbar c^3} - V \right) < 0$$



$$\frac{\dot{\phi}^2}{\hbar c^3 V} < 1$$

The potential energy must dominate the kinetic energy.

AS 4022 Cosmology

Scalar Field Dynamics

Equation of State: $w = p / \epsilon$

$$p_\phi = \frac{\dot{\phi}^2}{2\hbar c^3} - V(\phi)$$

$$\epsilon_\phi = \frac{\dot{\phi}^2}{2\hbar c^3} + V(\phi)$$

Evolution of energy density:

$$\epsilon \propto x^{3(1+w)} \quad dt = -dx / (xH)$$

$$\dot{\epsilon} = \frac{d\epsilon}{dx} \frac{dx}{dt} = \left(3(1+w) \frac{\epsilon}{x} \right) (-xH) = -3H(\epsilon + p)$$

Evolution of uniform scalar field:

$$\dot{\epsilon} = \frac{\dot{\phi}\ddot{\phi}}{\hbar c^3} + \frac{\partial V}{\partial \phi} \dot{\phi} = -3H(\epsilon + p) = -3H \frac{\dot{\phi}^2}{\hbar c^3}$$

$$\ddot{\phi} + 3H\dot{\phi} = -\hbar c^3 \frac{\partial V}{\partial \phi}$$

AS 4022 Cosmology


Scalar Field Dynamics

Hubble Drag:

Acceleration damped by expansion:

$$\ddot{\phi} + 3H\dot{\phi} - c^2 \nabla^2 \phi = -\hbar c^3 \frac{\partial V}{\partial \phi}$$

(Inflation makes spatial gradients small.)



"Slow-Roll" Approximation:

$$3H\dot{\phi} = -\hbar c^3 \frac{\partial V}{\partial \phi}$$

Terminal velocity:

$$\dot{\phi} \Rightarrow -\frac{\hbar c^3}{3H} \frac{\partial V}{\partial \phi}$$

"Like a snowflake falling"

Required for Inflation:

$$\frac{\dot{\phi}^2}{\hbar c^3 V} \Rightarrow \frac{\hbar c^3}{9H^2 V} \left(\frac{\partial V}{\partial \phi} \right)^2$$

Friedmann equation:


$$H^2 \approx \frac{8\pi G}{3} \frac{\epsilon}{c^2} \approx \frac{8\pi G}{3} \frac{V}{c^2}$$

$$= \frac{\hbar c^5}{24\pi G} \left(\frac{\partial V / \partial \phi}{V} \right)^2 \ll 1$$

Inflation requires a "very flat" potential.

AS 4022 Cosmology

Long Slow Roll Inflation Ending in a Hot Big Bang.



Long slow roll across the "plateau" (potential dominated) gives > 60 e-foldings of inflation (cooling, flattening).

Latent heat released during (kinetic energy dominated) rapid roll and damped oscillations at the end fills the universe with photons and particle-antiparticle pairs, launching the Hot Big Bang.

Initial quantum vacuum fluctuations mean different regions finish at slightly different times, giving the small (10^{-5}) temperature/density ripples we see on the CMB.

AS 4022 Cosmology

A Multi-verse? Chaotic Inflation?


Vacuum with quantum fluctuations.

Many bubbles of false vacuum inflate, with:
Different physical constants.
Different spatial dimensions.

ANTHROPIC PRINCIPLE:

Most bubbles quickly re-collapse or expand too fast to form stars or are otherwise unsuitable as habitats for Life.

Is our visible universe part of a big bubble lasting long enough, and with suitable physical laws, to allow beings like us to evolve?



AS 4022 Cosmology