

AS1001:Extra-Galactic Astronomy



Lecture 4: Galaxy Spectra

Galaxy Spectra

- The combined light from $\sim 10^{10}$ **stars** plus many **molecular clouds** and **star-forming regions**.
- The spectra tell us:
 - The galaxy's **velocity** (or redshift, hence distance)
 - The **mass** (from internal velocities)
 - The **star-formation rate** (emission lines)
 - The average **age** of the stellar population (blue/red)
- 3 Aspects of Spectra:
 - **Continuum**
 - **Absorption Lines**
 - **Emission Lines**

Galaxy Spectrum

Example galaxy spectrum:




Features:

Emission lines


Continuum

Absorption lines



The Continuum


- The combination of many Black-Body spectra from stars spanning a range in temperatures



- Red colour => lack of blue (hot young) stars
=> old stellar population
- Blue colour => ongoing star formation

Absorption Lines


- Atoms/Molecules in a star's atmosphere absorb light at specific wavelengths



- Cold gas in the interstellar medium (ISM) absorbs light at specific wavelengths.
- (ISM's cold dust absorbs ~ 1 mag / kpc)

Emission Lines

- Young stars are initially embedded in gas.
- Hot (high-mass) young stars ionise nearby gas.
- Gas emits at specific wavelengths as the free electrons recombine.



Hydrogen Energy Levels

Ionisation Potential

$$h \nu = 13.6 \text{ eV}$$

$$\lambda = 912 \text{ Angstroms}$$

UV photons ionise.


Recombinations then produce emission lines

$$\frac{1}{\lambda} = R \left(\frac{1}{n_l^2} - \frac{1}{n_u^2} \right)$$

$$R = 1.097 \times 10^7 \text{ m}^{-1}$$

$$= 1 / (912 \text{ \AA})$$

$$H\alpha : n_u = 3 \quad n_l = 2$$



Rydberg Formula for Hydrogen Lines


$$\frac{1}{\lambda} = R_{\infty} \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

Hydrogen Absorption Spectrum



**hot star
atmosphere**

Hydrogen Emission Spectrum



H II region

Orion Nebula

Hydrogen is ionized by photons
with $E > 13.6 \text{ eV}$ or $\lambda < 912 \text{ \AA}$.

$$1\text{eV} = 1.602\text{E-}19 \text{ J};$$


$$E = h \nu = h c / \lambda$$

Four bright O stars emit most of the
ionizing photons that produce the
Orion Nebula H II region

Neutral hydrogen: $\text{H}^0 = \text{H I}$

Ionized hydrogen: $\text{H}^+ = \text{H II}$

Similarly for other atoms/ions,
e.g. MgII , OIII , ...



HST View of Orion Nebula

Electrons recombine,
cascade thru energy levels,
emitting line photons.

Balmer lines in optical
Recombinations to $n = 2$


$H\alpha$: 6563 Å (red)

$O[III]$ 5007 Å (green)

$$\frac{1}{\lambda} = R \left(\frac{1}{n_l^2} - \frac{1}{n_u^2} \right)$$

$$R = 1.097 \times 10^7 \text{ m}^{-1}$$
$$= 1 / (912 \text{ Å})$$

$$H\alpha : n_u = 3 \quad n_l = 2$$






Orion Nebula HII region

H α map of Milky Way's Ionised Gas

Gas layer ~ 1 kpc thick. Ionized by O & B stars in the Galactic disc.



M101

Emission nebulae
(H II regions) along
the spiral arms.


Why are H II regions
pink?

Why are H II regions
along the spiral arms?



Model Spectra for a Starburst


A spectroscopic sequence



Absorption / Emission Lines

- Absorption Lines
 - hot: H, He cool: metals
 - Formed by atoms/molecules absorbing light
 - in stellar atmospheres
 - by cold gas in the ISM
- Implies
 - Metal lines from cool stars
=> old stellar population
=> old galaxy
- From
 - Ellipticals
 - Spiral Bulges
- Emission Lines
 - Gas ionised by UV photons from nearby O and B type stars
- Implies
 - Newly formed stars
=> star-forming ongoing
=> young galaxy
- From
 - Spiral Disks
 - Irregulars


Example Spectrum: Elliptical




The “4000Å-break”

- 4000 Angstrom = 400 nm
- Caused by:
 - **Absorption** (many overlapping lines with $\lambda < 4000\text{Å}$)
by **metals** in the atmospheres of cooler stars
 - **lack of hot blue stars** (type O,B)
- Hence:
 - Ellipticals \Rightarrow Strong 4000Å-Break
 - Spirals \Rightarrow Weak 4000Å-Break
 - Irregulars \Rightarrow Little or no 4000Å-Break

Example Spectrum: Spiral



Example Spectrum: Irregular



Radial Velocities


- Most galaxy spectra are **REDSHIFTED**.
- Observed wavelengths of spectral lines are longer than wavelengths measured for gasses in the lab.
- Interpret this as a DOPPLER shift.
- Most galaxies are moving away.
- $v > 0$: RECEDING
- $v < 0$: APPROACHING
- $z = \text{redshift}$ ($z = v / c$ for $z \ll 1$)
- **Distance** from Hubble's law: $d = v / H_0$ ($H_0 = 72 \text{ km/s/Mpc}$)

$$\frac{\lambda_{\text{OBSERVED}}}{\lambda_{\text{LAB}}} \equiv 1 + z = 1 + \frac{v}{c}$$

or

$$z = \frac{\Delta\lambda}{\lambda} = \frac{v}{c}$$

Example Radial Velocity



$$v = c \left(\frac{\lambda_{\text{OBS}} - \lambda_{\text{LAB}}}{\lambda_{\text{LAB}}} \right)$$
$$= (3 \times 10^5 \text{ km/s}) \left(\frac{4000 - 3727}{3727} \right) = 21,974 \text{ km/s}$$

$$v = c \left(\frac{467}{6563} \right) = 21,500 \text{ km/s}$$

GALAXY IS MOVING AWAY AT ABOUT 21,750 km/s

CLASS EXERCISE

- Work in groups of 2-3
- Collect an example spectrum
- Identify spectral features
- Measure the wavelengths of the spectral lines
- Calculate the radial velocity (km/s)
- Use Hubble's law ($d = H_0 v$) to find the distance
($H_0 = 72 \text{ km/s/Mpc}$)

Typical Spectral Features

- Absorption

- Ca II (K) = 3933.7 Å
- Ca II (H) = 3968.5 Å
- G-band = 4304.4 Å
- Mg I = 5175.3 Å
- Na I = 5894.0 Å

1 Angstrom = 0.1 nm = 10^{-10} m

- Emission

- [O II] = 3726.7 Å
- H δ = 4101.7 Å
- H γ = 4340.5 Å
- H β = 4861.3 Å
- [O III] = 4958.9 Å
- [O III] = 5006.8 Å
- H α = 6562.8 Å
- [S II] = 6716.0 Å

Brackets (e.g. [O III]) mean “forbidden lines”, emitted only at very low gas densities.